

<b>Title:</b>	Intraosseous needle insertion—paediatric guideline
<b>Version:</b>	<b>1</b>
<b>Supersedes:</b>	Not applicable
<b>Application:</b>	The guideline is intended for use by any hospital team caring for infants, children and young people under 16 years age across the Paediatric Critical Care Network in the North West & North Wales region.

<b>Originated /Modified By: Designation:</b>	Su Ying Ong, anaesthetic senior clinical fellow, NWTS / Alder Hey Children’s NHS Trust Isabel Wardach, senior clinical fellow, NWTS Kate Parkins, paediatric intensive care medicine consultant, NWTS Sophina Mahmood, paediatric pharmacist, PCC ODN & RMCH
<b>Ratified by:</b>	North West (England) & North Wales Paediatric Critical Care Operational Delivery Network, which includes multi-disciplinary clinical representation from all local and tertiary hospitals across the region.
<b>Date of Ratification:</b>	10.08.23
<b>Ratified by:</b>	RMCH (host trust for PCC ODN) policies and guidelines committee & Pharmacy and Medicines
<b>Date of Ratification:</b>	01.05.24

<b>Issue / Circulation Date:</b>	<b>1 Circulation date: 02.05.24</b>
<b>Circulated by:</b>	North West (England) & North Wales Paediatric Critical Care Operational Delivery Network,
<b>Dissemination and Implementation:</b>	Via networks December 2023
<b>Date placed on: NWTS &amp; PCC /LTV / SiC networks website</b>	03.05.24 NWTS Network website: May ‘24

<b>Planned Review Date:</b>	<b>3 years ie May 2027</b>
<b>Responsibility of:</b>	Clinical lead, North West & North Wales Paediatric Critical Care Network & NWTS guideline lead consultant

<b>EqIA Registration Number (RMCH):</b>	<b>2023-173</b>
---	-----------------

**1. Detail of Procedural Document**

Guidelines for insertion of paediatric intraosseous needle

**2. Equality Impact Assessment**

EqlA registration Number for RMCH:	<b>2023-173</b>
------------------------------------	-----------------

**3. Consultation, Approval and Ratification Process**

This guideline was developed with input from:

- North West (England) and North Wales Paediatric Transport Service (NWTS).
- North West and North Wales Paediatric Critical Care Operational Delivery Network
- Representatives from the District General Hospitals within network above.

These guidelines were circulated amongst the North West and North Wales Paediatric Critical Care Network for comments on 27.06.23

All comments received have been reviewed and appropriate amendments incorporated.

These guidelines were signed off by the Network Oversight Committee and Clinical Lead on 10.08.23

For ratification process for network guidelines see appendix 1.

**4. Disclaimer**

These clinical guidelines represent the views of the North West (England) and North Wales Paediatric Transport Service (NWTS) and the North West and North Wales Paediatric Critical Care Operational Delivery Network (PCCN). They have been produced after careful consideration of available evidence in conjunction with clinical expertise and experience.

It is intended that trusts within the Network will adopt this guideline and educational resource after review and ratification (including equality impact assessment) through their own clinical governance structures.

The guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient.

Clinical advice is always available from NWTS on a case by case basis.

Please feel free to **contact NWTS (01925 853 550)** regarding these documents if there are any queries

**Quick reference guide for paediatric intraosseous insertion**

**Patient requires vascular access**

Do you need immediate vascular access?

**NO**

Further discussion about need for peripheral or central access. IO access can be a reliable bridge until longer term vascular access can be established.

**YES**

Cardiac or respiratory arrest, impending arrest or shock or unstable dysrhythmia? **OR**  
Is peripheral access difficult &/or urgent treatment required eg anti-microbials/insulin in DKA/anti-convulsants in status epilepticus?

**YES**

Intraosseous (IO)

**NO**


Peripheral IV line

**Consider contraindications:**

- Fracture near or proximal to the insertion site
- Recent surgery to the limb or indwelling metal work
- IO insertion in the same site in the previous 48 hours
- Overlying infection or abscess
- Osteogenesis imperfecta; osteomyelitis
- Crush injury or ipsilateral vascular injury
- Landmarks not identifiable

**Needle selection:** check size of child & site of insertion, don't base it solely on weight


**Check the depth marker** to ensure at least **one black line is visible** above the skin **when needle has been pushed through skin & is resting on bone**



**Option 1: Proximal tibia**

**1. Position:**

Infant: flexed knee  
Child / Adolescent: straight leg



**2. Palpate tibial tuberosity** (bony thickness below patella)

**3. Insert 2-3cm below (or 2 FB) + 1 FB medial to tibia tuberosity at 90° to flat antero-medial surface of tibia**




**Option 2: Proximal humerus**

**1. Position:** Internal rotation of arm: bend arm at elbow & tuck hand behind pt's back  
**2. Palpate up** mid-shaft humerus towards humeral head to **locate surgical neck**




**3. Palpate greater tuberosity** (small bony protrusion directly above surgical neck)



**4. Insert at 45° to anterior plane (ie bed) into base of the greater tuberosity**

**Option 3: Distal femur**

- 1. Position:** leg outstretched
- 2. Palpate in mid-line**, 1-2 FB above & 1 FB medial to the superior border patella
- 3. Insert at approx. 15° cephalad** (towards head) to avoid growth plate and the tendon



## Step by Step Guide

1. Identify landmark, position patient and clean site with 2% chlorhexidine (if not allergic)

2. Place needle perpendicular (at 90° angle) to the bone (except for humerus at 45° angle to the bone)


3. **PUSH** (don't drill) needle through the skin & rest needle on bone. CHECK a minimum one black line is visible.

4. Start the drill (once through the skin) and gently drive into the bone without any additional force.  
Let the drill do the work

5. **STOP** when you feel loss of resistance or 'give' or 'pop'

6. CHECK that the needle is stable/solid in the bone—ie no 'wobble'

7. Unscrew and remove central stylet and dispose of sharp safely



8. Aspirate bone marrow to confirm location (NB not always possible) then send sample for culture and glucose if able (do NOT use blood gas machine as may block/damage analyser)

9. Secure with dressing (ideally ones with the EZ-IO set). Flush extension set with 0.9% sodium chloride then connect to IO

10. Confirm position by flushing with 5-10mL fluid via a syringe. Bolus will need more pressure than PVL.  
'No flush = no flow' ie NOT in correct position.


11. Check regularly for extravasation & monitor limb to check for dislodgement or compartment syndrome

12. Document accurately in patient's notes, including any sites where insertion failed.

**WARNING:** Avoid use of IO lidocaine as it has been associated with death or cardiac arrhythmias


**REMEMBER! AFTER INSERTION, CHECK:**

- Firmly seated needle (no wobble)
- Aspirate blood via a syringe (flash of blood)
- No leaking around site
- No sign of extravasation
- Secure eg using EZ stabiliser/sterile dressing or similar method
- EZ-connect/luer lock extension set
- Regularly for limb perfusion and any signs of extravasation or compartment syndrome (feel tissue / muscles surrounding insertion site and compare with opposite side. If it feels firmer /woody than the side without an IO the IO has tissue)
- Put pink IO name band on appropriate limb (leave in situ even if IO removed) to indicate which limb has had an IO



**INFUSIONS VIA IO**

- Attach a luer-lock extension line and then 3-tail extension line (see photo) to allow multiple compatible infusions to run via one IO
- IV fluids need to be infused under pressure or bolused using a 20 mL syringe.
- Gravity is insufficient to drive fluid through an IO
- All medications that can be given intravenously can be given intraosseously at the same doses.



**MONITOR / OBSERVATIONS**

- Check colour of the limb—should remain pink / healthy. Extravasation indicated if limb becomes pale / blue
- Presence of subcutaneous oedema, increasing limb size, tense muscle compartment (feels firmer or 'woody') compared to other limb, altered sensation, weak or absent distal pulses
- Position and fixation of the needle, patency of the IO, appearance of the insertion site (check for redness)
- Time elapsed since placement (ideally <24 hours)

**Potential Complications**

- Extravasation or subperiosteal infusion
- Dermal abrasion due to friction from the rotating plastic base surrounding the EZ-IO needle
- Compartment syndrome: rare but the smaller the patient the higher the risk
- Fracture or growth plate injury
- Osteomyelitis: very rare
- Fat embolus: rare

**REMOVAL**

- Ideally remove within 24 hours.
- Remove the EZ-connect extension set.
- Attach a clean 5 or 10mL luer lock syringe (acting as a handle/grip).
- Rotate the syringe clockwise.
- While rotating, gently pull the needle out, avoiding use of excessive force.
- Dispose of sharps safely.
- Apply pressure for a few minutes, if necessary then a small sterile dressing to the site.
- NB caution if coagulopathic—may need sustained pressure +/- platelets and clotting products.







**APPENDIX 1: INSERTION SITES**

**Needle selection:** check size of child & site of insertion.


**Don't base it solely on weight.**

**Check the depth marker AFTER the needle has been pushed through skin and is resting on bone (at 90° to surface of bone).**

When tip of needle is touching bone **STOP & CHECK** that at least **one black line (5 mm mark) is visible above the skin before starting to drill** ie drill/insert into bone, to ensure that the needle is long enough to reach the marrow.





Black 5 mm marker visible on IO needle after passing thru soft tissue & resting on bone/periosteum



Black 5 mm marker visible on IO needle after passing thru soft tissue & resting on bone/periosteum

**Option 1: Proximal tibia**

- POSITION:** Infant: flexed knee Child / Adolescent: straight leg
- PALPATE tibial tuberosity** = bony thickness below patella



NB Landmarks become more defined / easier to find as child gets older


**3. INSERTION:** if **tibial tuberosity CAN** be felt insert **1 finger breadth (FB) below & 1FB medial to tibia tuberosity**  
If **tibial tuberosity CANNOT** be felt, insert **2 FB below patella & 1 FB medial** along flat surface of tibia

**Aim to keep needle at 90° to the flat anteromedial surface of tibia**

**Neonate / infant**



**Older child**



**4. STOP** when you feel loss of resistance or 'give'

**Appendix 1 (continued)**


**Option 2: Proximal humerus**

**1. POSITION:** Internal rotation of arm 3 options:


**OPTION A:** Bend arm at elbow & place palm of hand on umbilicus, thumb up towards head



**OPTION B:** More ideally, tuck dorsal aspect of hand behind their back, resting against the hip (lying down: palm of hand against bed with thumb up towards head)




**OPTION C:** Place arm tight against body & rotate hand so palm is facing outwards, thumb pointing down to the floor






**2. PALPATE UP** mid-shaft humerus towards humeral head to **locate surgical neck** (narrower region).

**PALPATE greater tuberosity** (small bony protrusion directly above surgical neck)







**3. INSERT** at 45° to anterior plane (ie the patient's bed) into the base of the greater tuberosity





**Option 3: Distal femur**

1. **POSITION:** leg outstretched
2. **PALPATE** in **mid-line**, 1-2 FB above and 1 FB medial to the superior / upper border of patella
3. **INSERT:** angle needle at 90° to bone and approx. 15° cephalad (towards head) to avoid growth plate and the tendon






# RATIFICATION PATHWAY



**Manchester University**  
NHS Foundation Trust



**GUIDELINES:** [www.nwts.nhs.uk/clinicalguidelines](http://www.nwts.nhs.uk/clinicalguidelines)

**Crashcall link via NWTS website:** <https://www.nwts.nhs.uk/documentation/crashcall>

- for intubation drugs / sedation regime / inotropes etc

NWTS LocSIPPS / Checklists includes sizes of ETT, CVL & arterial lines

**EDUCATION:** [www.nwts.nhs.uk/education-website](http://www.nwts.nhs.uk/education-website)

Login details for education site are available from your nursing and medical paediatric critical care (PCC) operational delivery network (ODN) links

**Videos for IO insertion**

**ADULT intraosseous insertion:** <https://handbook.bcehs.ca/clinical-practice-guidelines/pr-clinical-procedure-guide/pr12-intraosseous-cannulation/>

**All Age Groups:** <https://www.teleflex.com/usa/en/clinical-resources/ez-io/>

**REFERENCES**

**FOR DRUG DOSES: British National Formulary for Children 2019-2020**

Advanced Life Support Group (ALSG). (2016). *Advanced Paediatric Life Support: A Practical Approach to Emergencies*. John Wiley & Sons Inc.

Nagler, J., & Krauss, B. (2011, February 24). Intraosseous Catheter Placement in Children. *The New England Journal of Medicine*. doi:10.1056/NEJMc0900916

Olson, D., Fesselet, J.-F., & Grouzard, V. (2018). Appendix 6. Intraosseous (IO) needle insertion in children. In *Management of a CHOLERA EPIDEMIC*. Médecins Sans Frontières. Retrieved from <https://medicalguidelines.msf.org/viewport/CHOL/english/appendix-6-intraosseous-io-needle-insertion-in-children-25297231.html#:~:text=The%20malleoli%20sites%20are%20approximately,where%20the%20bone%20is%20flat>.

Teleflex. (2022, February). Arrow EZ-IO Intraosseus Vascular Access: Care, Maintenance, and Removal. Retrieved from [https://www.teleflex.com/global/clinical-resources/documents/MCI-2019-0394\\_VA\\_DS\\_EZ-IO\\_Intraosseus\\_Vascular\\_Access\\_LR.pdf](https://www.teleflex.com/global/clinical-resources/documents/MCI-2019-0394_VA_DS_EZ-IO_Intraosseus_Vascular_Access_LR.pdf)

The Royal Children's Hospital Melbourne. (2022, February). Intraosseous access - Clinical Practice Guideline. Retrieved from [https://www.rch.org.au/clinicalguide/guideline\\_index/Intraosseous\\_access/](https://www.rch.org.au/clinicalguide/guideline_index/Intraosseous_access/)

Gavin Owen and Owen Keane. Intraosseous Access, Don't Forget The Bubbles, 2020. Available at <https://dontforgetthebubbles.com/author/gavinowen/>

**Contact numbers:**

Regional Paediatric Intensive Care Unit Alder Hey Childrens Hospital 0151 252 5241

Regional Paediatric Intensive Care Unit Royal Manchester Childrens Hospital 0161 701 8000

NWTS (North West & North Wales Paediatric Transport Service) 01925 853 550

**Guideline authors:**

Dr Su Ying Ong, senior anaesthetic fellow, AHCH

Dr Isabel Wardach, senior clinical fellow, NWTS

Dr Kate Parkins, PICM consultant, NWTS

**Consulted parties:**

North West & North Wales Paediatric Transport Service (NWTS)

North West and North Wales Paediatric Critical Care Network

PICU, Royal Manchester Children's Hospital

PICU, Alder Hey Children's Hospital

**GUIDELINE CONTACT POINT:** [Kate.parkins@nwts.nhs.uk](mailto:kate.parkins@nwts.nhs.uk) & [Nicola.longden@mft.nhs.uk](mailto:Nicola.longden@mft.nhs.uk)

Please visit NWTS website for the most up to date version of this guideline: [www.nwts.nhs.uk/guidelines](http://www.nwts.nhs.uk/guidelines)

**Date of Approval by Paediatric Critical Care ODN: 10.08.23**

**Date of Approval by Host Trust (RMCH/MFT): 01.05.24**

**Date of Review: May 2027**